


AMENDMENTS TO THE CLAIMS

Claims 1-7 (canceled)

8. (Currently amended) A method of prevention or treatment of a disorder in a mammal, the said disorder being selected from the group consisting of toxic effects of TNF- α , alcohol-induced hepatitis, and cachexia, comprising administering to said mammal an effective amount of a pteridine derivative having the structural formula (I):



wherein X represents an oxygen atom or a group with the formula S(O)_m wherein m is an integer from 0 to 2, or a group with the formula NZ and wherein:

- R₁ is selected from the group consisting of methyl, ethyl, isopropyl and pentyl;
- Z is a group independently defined as R₁ or Z is hydrogen or the group NZ together with R₁ is either hydroxylamino or an optionally substituted heterocyclic group containing at least one nitrogen atom;
- R₂ is selected from the group consisting of amino; acylamino;
- R₄ is an atom or a group selected from the group consisting of hydrogen; halogen; C₁₋₇ alkyl; C₂₋₇ alkenyl; C₂₋₇ alkynyl; halo C₁₋₇ alkyl; carboxy C₁₋₇ alkyl; acetoxy C₁₋₇ alkyl; carboxyaryl; C₁₋₇ alkoxy; C₃₋₁₀ cycloalkoxy; aryloxy; arylalkyloxy; oxyheterocyclic; heterocyclic-substituted alkyloxy; thio C₁₋₇ alkyl; thio C₃₋₁₀ cycloalkyl; thioaryl; thioheterocyclic; arylalkylthio; heterocyclic-substituted alkylthio; amino; hydroxylamino; mercapto-amino; acylamino; thioacylamino; alkoxyamino; thioalkylamino; acetal; thioacetal; carboxylic acid; carboxylic acid esters, thioesters, halides, anhydrides, amides and thioamides; thiocarboxylic acid; thiocarboxylic acid esters, thioesters, halides, anhydrides, amides and thioamides; hydroxyl; sulfhydryl; nitro; cyano; carbamoyl; thiocarbamoyl, ureido; thio-ureido;

alkylamino; cycloalkylamino; alkenylamino; cycloalkenylamino; alkynyl-amino; arylamino; arylalkylamino; hydroxyalkylamino; mercapto-alkylamino; heterocyclic amino; heterocyclic-substituted alkylamino; oximino; alkyloximino; hydrazino; alkylhydrazino; phenylhydrazino; cysteinyl acid, esters, thioesters, halides, anhydrides, amides and thioamides thereof; aryl groups optionally substituted with one or more substituents selected from the group consisting of halogen, C₁₋₇ alkyl, C₁₋₇ alkoxy; optionally substituted heterocyclic radicals; aromatic or heterocyclic substituents substituted with an aliphatic spacer between the pteridine ring and the aromatic or heterocyclic substituent, whereby said aliphatic spacer is a branched or straight, saturated or unsaturated aliphatic chain of 1 to 4 carbon atoms; branched or straight, saturated or unsaturated aliphatic chains of 1 to 7 carbon atoms; and

- R₃ is an atom or a group defined as R₄, or R₃ together with R₄ forms a homocyclic or heterocyclic radical;

and/or being a pharmaceutically acceptable addition salt thereof and/or a stereoisomer thereof and/or a mono- or a di-N-oxide thereof and/or a solvate and/or a dihydro- or tetrahydropteridine derivative thereof.

9. (Previously presented) A method according to claim 8, wherein R₄ is hydrogen or methoxy.

10. (Previously presented) A method according to claim 8, wherein R₄ is hydrogen and wherein R₂ is amino.

11. (Previously presented) A method according to claim 8, wherein R₃ is 3-thienyl, 2-thienyl or a phenyl group with one or more substituents.

12. (Previously presented) A method according to claim 8, wherein R₂ is amino and wherein R₃ is 3-thienyl, 2-thienyl or a phenyl group with one or more substituents.
13. (Previously presented) A method according to claim 8, wherein R₃ is 3-thienyl, 2-thienyl or a phenyl group with one or more substituents, and wherein R₄ is hydrogen.
14. (Previously presented) A method according to claim 8, wherein R₂ is amino and wherein R₃ is 3-thienyl, 2-thienyl or a phenyl group with one or more substituents, and wherein R₄ is hydrogen.
15. (Previously presented) A method according to claim 8, wherein R₃ is a phenyl group with one or more substituents each independently selected from the group consisting of fluoro, methoxy, ethoxy, trifluoromethyl, dimethylamino, chloro, cyano, methyl, ethyl, carboxymethyl, methylthio, dimethylcarboxamido, diethylcarboxamido and methylcarboxylate.
16. (Previously presented) A method according to claim 8, wherein R₄ is hydrogen and wherein R₃ is a phenyl group with one or more substituents each independently selected from the group consisting of fluoro, methoxy, ethoxy, trifluoromethyl, dimethylamino, chloro, cyano, methyl, ethyl, carboxymethyl, methylthio, dimethylcarboxamido, diethylcarboxamido and methylcarboxylate.
17. (Previously presented) A method according to claim 8, wherein R₂ is amino, wherein R₄ is hydrogen and wherein R₃ is a phenyl group with one or more substituents each independently selected from the group consisting of fluoro, methoxy, ethoxy,

trifluoromethyl, dimethylamino, chloro, cyano, methyl, ethyl, carboxymethyl, methylthio, dimethylcarboxamido, diethylcarboxamido and methylcarboxylate.

18. (Previously presented) A method according to claim 8, wherein R₃ is selected from the group consisting of 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 3,4-difluorophenyl, 4-dimethylaminophenyl, 4-trifluoromethylphenyl, 3,4-dichlorophenyl, 4-cyanophenyl, 4-ethoxyphenyl, 4-fluorophenyl, 4-ethylphenyl, 3-fluoro-4-methylphenyl, 3-methyl-4-methoxyphenyl, 3,4-dimethylphenyl, 3-chloro-4-trifluoromethylphenyl, 4-acetylphenyl, 3,4-dimethoxyphenyl, styryl, 4-tolyl, 4-chlorophenyl and 3,4,5-trimethoxyphenyl.
19. (Previously presented) A method according to claim 8, wherein R₂ is amino, and wherein R₃ is selected from the group consisting of 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 3,4-difluorophenyl, 4-dimethylaminophenyl, 4-trifluoro-methylphenyl, 3,4-dichlorophenyl, 4-cyanophenyl, 4-ethoxyphenyl, 4-fluorophenyl, 4-ethylphenyl, 3-fluoro-4-methylphenyl, 3-methyl-4-methoxyphenyl, 3,4-dimethylphenyl, 3-chloro-4-trifluoromethylphenyl, 4-acetylphenyl, 3,4-dimethoxyphenyl, styryl, 4-tolyl, 4-chlorophenyl and 3,4,5-trimethoxyphenyl.
20. (Previously presented) A method according to claim 8, wherein R₄ is hydrogen, and wherein R₃ is selected from the group consisting of 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 3,4-difluorophenyl, 4-dimethylaminophenyl, 4-trifluoro-methylphenyl, 3,4-dichlorophenyl, 4-cyanophenyl, 4-ethoxyphenyl, 4-fluorophenyl, 4-ethylphenyl, 3-fluoro-4-methylphenyl, 3-methyl-4-methoxyphenyl, 3,4-dimethylphenyl, 3-chloro-4-trifluoromethylphenyl, 4-acetylphenyl, 3,4-dimethoxyphenyl, styryl, 4-tolyl, 4-chlorophenyl and 3,4,5-trimethoxyphenyl.

21. (Previously presented) A method according to claim 8, wherein R₂ is amino, wherein R₄ is hydrogen, and wherein R₃ is selected from the group consisting of 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 3,4-difluorophenyl, 4-dimethylaminophenyl, 4-trifluoromethylphenyl, 3,4-dichlorophenyl, 4-cyanophenyl, 4-ethoxyphenyl, 4-fluorophenyl, 4-ethylphenyl, 3-fluoro-4-methylphenyl, 3-methyl-4-methoxyphenyl, 3,4-dimethylphenyl, 3-chloro-4-trifluoromethylphenyl, 4-acetylphenyl, 3,4-dimethoxyphenyl, styryl, 4-tolyl, 4-chlorophenyl and 3,4,5-trimethoxyphenyl.

22. (Previously presented) A method according to claim 8, wherein:

- X is NZ,
- Z is selected from the group consisting of hydrogen, methyl, ethyl, n-propyl and benzyl; and
- R₁ is selected from the group consisting of methyl, ethyl, n-propyl and benzyl.

23. (Previously presented) A method according to claim 8, wherein X is NZ and wherein the group NZ together with R₁ is selected from the group consisting of tetrahydropyridinyl, hydroxylamino, morpholinyl, piperidinyl, piperazinyl, 1,2,4-triazolyl and N-methylpiperazinyl.

24. (Previously presented) A method according to claim 8, wherein R₂ is amino, wherein R₄ is hydrogen, wherein X is NZ and wherein the group NZ together with R₁ is selected from the group consisting of tetrahydropyridinyl, hydroxylamino, morpholinyl, piperidinyl, piperazinyl, 1,2,4-triazolyl and N-methyl-piperazinyl.

25. (Previously presented) A method according to claim 8, wherein the pteridine derivative is administered in combination with one or more pharmaceutically acceptable carriers or excipients.
26. (Previously presented) A method according to claim 8, wherein the pteridine derivative is in the form of a hydrate or a solvate with an organic solvent selected from the group consisting of alcohols, ketones and esters.
27. (Previously presented) A method according to claim 8, wherein the pteridine derivative is a compound selected from the group consisting of:
- 2-amino-4-ethoxypteridine
 - 2-amino-4-ethoxy-6-chloro-pteridine
 - 2-amino-4-ethoxy-6-(4-methoxyphenyl)-pteridine
 - 2-amino-4-ethoxy-6-(2-methoxyphenyl)-pteridine
 - 2-amino-4-ethoxy-6-(3-methoxyphenyl)-pteridine
 - 2-amino-4-ethoxy-6-(3,4-difluorophenyl)-pteridine
 - 2-amino-4-ethoxy-6-(4-dimethylaminophenyl)-pteridine
 - 2-amino-4-ethoxy-6-(4-trifluoromethylphenyl)-pteridine
 - 2-amino-4-ethoxy-6-(2-thienyl)-pteridine
 - 2-amino-4-ethoxy-6-(3-thienyl)-pteridine
 - 2-amino-4-ethoxy-6-(3,4-dichlorophenyl)-pteridine
 - 2-amino-4-ethoxy-6-(4-cyanophenyl)-pteridine
 - 2-amino-4-ethoxy-6-(4-ethoxyphenyl)-pteridine
 - 2-amino-4-ethoxy-6-(4-fluorophenyl)-pteridine
 - 2-amino-4-ethoxy-6-(4-ethylphenyl)-pteridine
 - 2-amino-4-ethoxy-6-(4-acetylphenyl)-pteridine

- 2-amino-4-ethoxy-6-(3-fluoro-4-methylphenyl)-pteridine
- 2-amino-4-ethoxy-6-(4-methylthiophenyl)-pteridine
- 2-amino-4-ethoxy-6-(4-N,N-dimethylbenzamido)-pteridine
- 2-amino-4-isopropoxypteridine
- 2-amino-4-isopropoxy-6-chloropteridine
- 2-amino-4-isopropoxy-6-(3-methyl-4-methoxyphenyl)-pteridine
- 2-amino-4-isopropoxy-6-(3,4-dimethylphenyl)-pteridine
- 2-amino-4-isopropoxy-6-(3-chloro-4-trifluoromethylphenyl)-pteridine
- 2-amino-4-isopropoxy-6-(3-chlorol-4-fluorophenyl)-pteridine
- 2-amino-4-isopropoxy-6-(4-N,N-diethylbenzamido)-pteridine
- 2-amino-4-isopropoxy-6-(4-trifluoromethylphenyl)-pteridine
- 2-amino-4-isopropoxy-6-(3,4-difluorophenyl)-pteridine
- 2-amino-4-isopropoxy-6-(4-methoxyphenyl)-pteridine
- 2-amino-4-isopropoxy-6-(4-ethoxyphenyl)-pteridine
- 2-amino-4-isopropoxy-6-(4-N,N-dimethylbenzamido)-pteridine
- 2-amino-4-isopropoxy-6-(3-thienyl)-pteridine
- 2-amino-4-isopropoxy-6-(4-cyanophenyl)-pteridine
- 2-amino-4-isopropoxy-6-(4-benzoic acid methyl ester)-pteridine
- 2-amino-4-isopropoxy-6-(4-acetylphenyl)-pteridine
- 2-amino-4-isopropoxy-6-(3,4-dimethoxyphenyl)-pteridine
- 2-amino-4-ethylthio-6-(3,4-dimethoxyphenyl)-pteridine
- 2-amino-4-isopropylthio-6-(3,4-dimethoxyphenyl)-pteridine
- 2-amino-4-pentoxy-6-styrylpteridine,
- 2-amino-4-n-pentoxy-6-(1,2-dibromo-2-phenylethyl)-pteridine,
- 2-amino-4-methoxy-6-styryl-7-methoxypteridine,
- 2-amino-4-dimethylamino-6-phenylpteridine,
- 2-amino-4-dimethylamino-6-(4-tolyl)pteridine,

- 2-amino-4-dimethylamino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-diethylamino-6-phenylpteridine,
- 2-amino-4-diethylamino-6-(4-chlorophenyl)pteridine,
- 2-amino-4-diethylamino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-diethylamino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-dipropylamino-6-phenylpteridine,
- 2-amino-4-dipropylamino-6-(4-chlorophenyl)pteridine,
- 2-amino-4-dipropylamino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-dipropylamino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-morpholino-6-phenylpteridine,
- 2-amino-4-morpholino-6-(4-chlorophenyl)pteridine,
- 2-amino-4-morpholino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-morpholino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-piperidino-6-phenylpteridine,
- 2-amino-4-piperidino-6-(4-chlorophenyl) pteridine,
- 2-amino-4-piperidino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-piperidino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-N-methylpiperazino-6-phenylpteridine,
- 2-amino-4-N-methylpiperazino-6-(4-chlorophenyl)pteridine,
- 2-amino-4-N-methylpiperazino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-methylpiperazino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-pyrrolidino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-piperazino-6-phenylpteridine,
- 2-amino-4-piperazino-6-(4-chlorophenyl)pteridine,
- 2-amino-4-piperazino-6-(4-methoxyphenyl)pteridine,
- 2-amino-4-piperazino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-morpholino-6-(3,4,5-trimethoxyphenyl)pteridine,

- 2-amino-4-morpholino-6-(3,4-formylidene-3,4-dihydroxyphenyl)pteridine,
- 2-amino-4-dimethylamino-6-(3,4-formylidene-3,4-dihydroxyphenyl) pteridine,
- 2-amino-4-pyrrolidino-6-(3,4,dimethoxyphenyl)pteridine,
- 2-amino-4-dimethylamino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-dimethylamino-6-methylpteridine,
- 2-amino-4-ethoxy-6-phenylpteridine,
- 2-amino-4-propylamino-6-phenylpteridine,
- 2-amino-4-propylamino-6-(3,4-dimethoxyphenyl)pteridine,
- 2-acetamido-4-isopropoxy-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-ethoxy-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-(1,2,3,6-tetrahydropyridinyl)-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-ethoxy-pteridine,
- 2-amino-4-ethoxypteridine-N⁸-oxide,
- 2-amino-4-isopropoxypertidine-N⁸-oxide,
- 2-amino-6-chloro-4-ethoxypteridine,
- 2-amino-6-chloro-4-isopropoxypertidine,
- 2-amino-6-(p-methoxyphenyl)-4-ethoxy-pteridine;
- 2-amino-6-(o-methoxyphenyl)-4-ethoxy-pteridine;
- 2-amino-6-(m-methoxyphenyl)-4-ethoxy-pteridine;
- 2-amino-6-(3,4-difluorophenyl)-4-ethoxy-pteridine;
- 2-amino-6-(p-dimethylaminophenyl)-4-ethoxy-pteridine;
- 2-amino-6-(p-trifluoromethylphenyl)-4-ethoxy-pteridine;
- 2-amino-6-(2-thienyl)-4-ethoxy-pteridine;
- 2-amino-6-(3-thienyl)-4-ethoxy-pteridine;
- 2-amino-6-(3,4-dichlorophenyl)-4-ethoxy-pteridine;
- 2-amino-6-(p-cyanophenyl)-4-ethoxy-pteridine;
- 2-amino-6-(p-ethoxyphenyl)-4-ethoxy-pteridine;

- 2-amino-6-(p-fluorophenyl)-4-ethoxy-pteridine;
- 2-amino-6-(p-ethylphenyl)-4-ethoxy-pteridine;
- 2-amino-6-(p-acetylphenyl)-4-ethoxy-pteridine;
- 2-amino-6-(3-methyl-4-fluorophenyl)-4-ethoxy-pteridine;
- 2-amino-6-(p-thiomethylphenyl)-4-ethoxy-pteridine;
- 2-amino-6-(p-N,N-dimethylbenzamido)-4-ethoxy-pteridine;
- 2-amino-6-(3,4-dimethoxyphenyl)-4-ethoxy-pteridine,
- 2-amino-6-(3-methyl-4-methoxyphenyl)-4-isopropoxyppteridine;
- 2-amino-6-(3,4-dimethylphenyl)-4-isopropoxyppteridine;
- 2-amino-6-(3-chloro-4-trifluoromethylphenyl)-4-isopropoxyppteridine;
- 2-amino-6-(3-chloro-4-fluorophenyl)-4-isopropoxyppteridine;
- 2-amino-6-(p-N,N-diethylbenzamido)-4-isopropoxyppteridine;
- 2-amino-6-(p-trifluoromethylphenyl)-4-isopropoxyppteridine;
- 2-amino-6-(3,4-difluorophenyl)-4-isopropoxyppteridine;
- 2-amino-6-(p-methoxyphenyl)-4-isopropoxyppteridine;
- 2-amino-6-(p-ethoxyphenyl)-4-isopropoxyppteridine;
- 2-amino-6-(p-dimethylbenzamido)-4-isopropoxyppteridine;
- 2-amino-6-(3-thienyl)-4-isopropoxyppteridine;
- 2-amino-6-(p-cyanophenyl)-4-isopropoxyppteridine;
- 2-amino-6-(p-benzoic acid methyl ester)-4-isopropoxyppteridine;
- 2-amino-6-(p-acetylphenyl)-4-isopropoxyppteridine;
- 2-amino-6-(3,4-dimethoxyphenyl)-4-isopropoxyppteridine,
- 2-amino-4-mercaptopethyl-6-(3,4-dimethoxyphenyl) pteridine;
- 2-amino-4-mercaptopisopropyl-6-(3,4-dimethoxyphenyl) pteridine,
- 2-acetylamino-4-(1,2,4-triazolyl)-6-(p-methoxyphenyl) pteridine,
- 2-acetylamino-4-(1,2,4-triazolyl)-7-(p-methoxyphenyl)pteridine,
- 2-amino-4-isopropoxy-7-(p-methoxyphenyl) pteridine,

- 2-amino-4-isopropoxy-7-(3,4-dimethoxyphenyl) pteridine,
- 2-amino-4-ethoxy-7-(3,4-dimethoxyphenyl) pteridine,
- 2-amino-4-methoxy-7-(3,4-dimethoxyphenyl) pteridine,
- 2-amino-4-(1,2,3,6-tetrahydropyridinyl)-6-(3,4-dimethoxyphenyl)pteridine,
- 2-amino-4-(diethanolamino)-6-[[3,4-(dimethoxyphenyl)]pteridine,
- 2-amino-4-thiomorpholino-6-[[3,4-(dimethoxyphenyl)]pteridine,
- 2-amino-4-morpholino-6-(4-acetanilide) pteridine,
- 2-amino-4-morpholino-6-(3-acetanilide) pteridine,
- 2-amino-4-morpholino-6-(4-aminophenyl) pteridine,
- 2-amino-4-morpholino-6-(3-aminophenyl) pteridine,
- 2-amino-4-morpholino-6-(4-benzoylaminophenyl) pteridine;
- 2-amino-4-morpholino-6-(4-phenoxyacetylaminophenyl) pteridine;
- 2-amino-4-morpholino-6-(4-propionylaminophenyl) pteridine;
- 2-amino-4-morpholino-6-(4-furoylaminophenyl) pteridine;
- 2-amino-4-morpholino-6-(4-cyclohexanoylaminophenyl) pteridine;
- 2-amino-4-morpholino-6-[4-(4-chlorobenzoyl)aminophenyl] pteridine;
- 2-amino-4-morpholino-6-(4-benzyloxyacetylaminophenyl) pteridine,
- 2-amino-4-morpholino-6-(4-isonicotinoylaminophenyl) pteridine;
- 2-amino-4-morpholino-6-(4-naphtoylaminophenyl) pteridine;
- 2-amino-4-morpholino-6-(4-methylsulfonylaminophenyl) pteridine;
- 2-amino-4-morpholino-6-(4-ethylsuccinylaminophenyl) pteridine;
- 2-amino-4-morpholino-6-[4-(4-methylbenzoate)aminophenyl) pteridine;
- 2-amino-4-morpholino-6-(3-benzoylaminophenyl) pteridine;
- 2-amino-4-morpholino-6-(3-benzensulfonylaminophenyl) pteridine;
- 2-amino-4-morpholino-6-(3-phenoxyacetylaminophenyl) pteridine;
- 2-amino-4-morpholino-6-(3-isonicotinoylaminophenyl) pteridine;
- 2-amino-4-morpholino-6-(3-cyclohexanoylaminophenyl) pteridine;

- 2-amino-4-morpholino-6-[3-(4-methylbenzoate)aminophenyl] pteridine;
- 2-amino-4-morpholino-6-(3-ethylsuccinylaminophenyl) pteridine;
- 2-amino-4-morpholino-6-(3-ethylmalonylaminophenyl) pteridine;
- 2-amino-4-morpholino-6-(3-benzyloxyacetylaminophenyl) pteridine;
- 2-amino-4-morpholino-6-(3-ethylsulfonylaminophenyl)pteridine,
- 2-amino-4-morpholino-6-[3-Boc-(L)-phenylalanine-aminophenyl] pteridine;
- 2-amino-4-morpholino-6-[3-Boc-(D)-phenylalanine-aminophenyl] pteridine;
- 2-amino-4-morpholino-6-[3-Boc-(L)-tryptophane-aminophenyl] pteridine;
- 2-amino-4-morpholino-6-[3-Boc-(D)-tryptophane-aminophenyl] pteridine,
- 2-amino-4-morpholino-6-(4-hydroxyphenyl) pteridine,
- 2-amino-4-morpholino-6-(4-ethoxyphenyl) pteridine;
- 2-amino-4-morpholino-6-(4-benzyloxyphenyl) pteridine;
- 2-amino-4-morpholino-6-(4-(phenethyloxy)-phenyl) pteridine;
- 2-amino-4-morpholino-6-(4-phenoxy-butyronitrile) pteridine;
- 2-amino-4-morpholino-6-(4-propoxy-phenyl) pteridine;
- 2-amino-4-morpholino-6-(4-phenoxy-butyric acid ethyl ester) pteridine;
- 2-amino-4-morpholino-6-(4-phenoxy-acetic acid ethyl ester) pteridine
- 2-amino-4-morpholino-6-(4-(2-methoxyethoxy)-phenyl) pteridine; and
- 2-amino-4-morpholino-6-(4-butoxy-phenyl)-pteridine.